

ON A THEOREM OF HARER

ROBERT TREGER

INTRODUCTION

Let $\mathbf{M}_g$ be the coarse moduli space of smooth projective complex curves of genus g . Let $\text{Pic}(\mathbf{M}_g)$ denote the Picard group of line bundles on $\mathbf{M}_g$. According to Mumford, $\text{Pic}(\mathbf{M}_g) \otimes \mathbb{Q} \simeq H^2(\mathbf{M}_g, \mathbb{Q})$ for $g \geq 2$ [9].

The purpose of this note is to give another proof of the following fundamental theorem of Harer [4].

Theorem 1. $\text{rk Pic}(\mathbf{M}_g) \otimes \mathbb{Q} = 1$ for $g \geq 3$.

We will deduce Theorem 1 from the Harer stability theorem [5, 6, 8].

Theorem 2. $H^2(\mathbf{M}_g, \mathbb{Q}) \simeq H^2(\mathbf{M}_{g+1}, \mathbb{Q})$ for $g \geq 6$.

Note that Theorem 2 is similar to the corresponding stability results for arithmetic groups.

PROOF OF THEOREM 1

To deduce Theorem 1 from Theorem 2, it suffices to show that $\text{rk Pic}(\mathbf{M}_g) \otimes \mathbb{Q} = 1$ for $g = 3, 4, 5, 6$. Let $V_{n,d} \subset \mathbb{P}^N$ ($N = n(n+3)/2$) be the variety of irreducible complex plane curves of degree n with d nodes and no other singularities. Let $V(n, g) \subset \mathbb{P}^N$ ($N = n(n+3)/2$) be the variety of irreducible complex plane curves of degree n and genus $g = (n-1)(n-2)/2 - d$. Clearly $V_{n,d} \subset V(n, g)$. Let

$$\Sigma_{n,d} \subset \mathbb{P}^N \times \text{Sym}^d(\mathbb{P}^2)$$

be the *closure* of the locus of pairs $(E, \sum_{i=1}^d R_i)$, where E is an irreducible nodal curve and $R_1, \dots, R_d$ are its nodes, and $\pi_{n,d}$ the projection to the symmetric product. We fix a general point P and a general line L in $\mathbb{P}^2$, and consider the following two elements in $\text{Pic}(V_{n,d})$: (CP) = the divisor class of curves containing the point P , and (NL) = the divisor class of curves with a node located somewhere on L .

If $g \geq 3(g+2-n)$ then the natural map $V(n, g) \rightarrow \mathbf{M}_g$ is a *surjective* morphism [3, p. 358]. Precisely, for any smooth curve C of genus g , there is a line bundle $\mathcal{L}$ on C of degree n such that $h^0(C, \mathcal{L}) \geq 3$. Furthermore, Harris observed that $\text{rk Pic}(\mathbf{M}_g) \otimes \mathbb{Q} = 1$ provided one can show that $\text{Pic}(V_{n,d})$ is a torsion group, where $d = (n-1)(n-2)/2 - g$ [7]. Thus Theorem 1 follows from the

1991 *Mathematics Subject Classification*. Primary 14H10.

Key words and phrases. Moduli of curves, plane curves, Picard group.

Lemma. *Let n and d be a pair of positive integers such that one of the following conditions holds: i) $n \gg d$ or ii) $(n, d) = (5, 3), (6, 6), (6, 5),$ or $(6, 4)$. Then $\text{Pic}(V_{n,d})$ is a torsion group generated by (CP) and (NL) .*

Proof of Lemma. We will prove the lemma for $(n, d) = (6, 6)$, hence $g = 4$. The remaining cases are similar only easier. Given *any* 6 distinct points $q_1, \dots, q_6$ in $\mathbb{P}^2$, no four on a line, then a general curve C of the linear system $\mathcal{L}$ of curves of degree 6 with assigned singularities at $q_1, \dots, q_6$ has 6 nodes and no other singularities. We will calculate $\dim \mathcal{L}$ using Castelnuovo's method (see [1, Book IV, Chap. I, Sect. 3]).

The characteristic series determined on C by $\mathcal{L}$ is cut out by curves of $\mathcal{L}$. Clearly one can find two curves in $\mathcal{L}$, say C_1 and C_2 , such that $(C_1 \cdot C_2)_{q_i} = 4$ for $i = 1, \dots, 6$. It follows that the degree of the characteristic series equals $(\deg C)^2 - 6 \cdot 4 > 6 = 2g - 2$. Hence this series is non-special, the superabundance of $\mathcal{L}$ vanishes, and $\dim \mathcal{L} = 6(6 + 3)/3 - 6 \cdot 3$.

Let S be a closed subset of $\text{Sym}^6(\mathbb{P}^2)$ which is a union of the singular locus and the cycles with at least 4 points on a line. Clearly $\text{codim } S = 2$ and $\text{Pic}(\text{Sym}^6(\mathbb{P}^2) \setminus S) = \text{Pic}(\text{Sym}^6(\mathbb{P}^2)) = \mathbb{Z}$. We set

$$\Sigma = \pi_{6,6}^{-1}(\text{Sym}^6(\mathbb{P}^2) \setminus S) \cap \Sigma_{6,6}.$$

By the above discussion, all the fibers of $\pi_{6,6}|_\Sigma$ have the same dimension and $\pi_{6,6}|_\Sigma: \Sigma \rightarrow \text{Sym}^6(\mathbb{P}^2) \setminus S$ is a projective bundle. Thus $\text{Pic}(\Sigma) = \mathbb{Z}^2$ and its generators correspond to the divisor classes (CP) and (NL) coming from the fiber and the base of the fibration, respectively.

We consider two divisors on Σ : (i) the closure of the locus of irreducible curves with $d - 1$ assigned nodes and one assigned cusp, and (ii) the closure of the locus of nodal curves with d assigned nodes and one unassigned node. The corresponding elements of $\text{Pic}(\Sigma) \otimes \mathbb{Q}$ are linearly independent [2, Sect. 2]. There is a natural open immersion $V_{6,6} \subset \Sigma$. Hence we get an exact sequence

$$\mathbb{Z}^r \rightarrow \text{Pic}(\Sigma) \rightarrow \text{Pic}(V_{6,6}) \rightarrow 0$$

where $\mathbb{Z}^r$ is generated by the irreducible components of $\Sigma \setminus V_{6,6}$ of codimension one in Σ . Therefore $\text{Pic}(V_{6,6})$ is torsion. This proves the lemma and the theorem.

REFERENCES

1. J. L. Coolidge, *Algebraic Plane Curves*, Oxford University Press, Oxford, 1931.
2. S. Diaz and J. Harris, *Geometry of the Severi variety II : Independence of the divisor classes and examples*, Algebraic Geometry. Sundance 1986, Lect. Notes Math. **1311**, Springer-Verlag, Berlin Heidelberg New York Tokyo, 1988, pp. 23-50.
3. P. Griffiths and J. Harris, *Principles of Algebraic Geometry*, John Wiley & Sons, New York, 1978.
4. J. Harer, *The second homology group of the mapping class group of an orientable surface*, Invent. Math. **72** (1982), 221-239.
5. J. Harer, *Stability of the homology of the mapping class group of orientable surfaces*, Ann. Math. **121** (1985), 215-249.
6. J. Harer, *The cohomology of the moduli space of curves*, Theory of Moduli (E. Sernesi, ed.), Lect. Notes Math. **1337**, Springer-Verlag, Berlin Heidelberg New York, 1985, pp. 138-221.
7. J. Harris, *Personal communication*.

8. N. V. Ivanov, *On the homology stability for the Teichmüller modular groups: closed surfaces and twisted coefficients*, Mapping Class Groups and Moduli Spaces of Riemann Surfaces (C. - F. Bökinger and R. M. Hain, eds.), Contemp. Math. **150**, Amer. Math. Society, Providence, RI, 1994, pp. 149-194.
9. D. Mumford, *Abelian quotients of the Teichmüller modular group*, J. d'Analyse Math. **18** (1967), 227-244.

AG&N, PRINCETON, NEW JERSEY
E-mail address: `tregrob@aol.com`